

No.	NAME	Age	Admitted	Died	Under the care of	Examination, at what time after death	No. in Reg ^t 1877	NATURE OF DISEASE
20	Thomas Hindley	50	18 th Jan 77	18 th Jan	W ^c Holmes	26 hours	82	Fractured ribs. Haemothorax. Luxation of spine

MORBID APPEARANCES

5 f. 8 Body well nourished, full habit. No visible trace of external injury, with the exception of a little marbling of the skin at precordial region and some bulging of the R side of thorax posteriorly.

Hair light brown

Lungs. The R. pleura contained about 2 pints of fluid blood. The L. pleura rather more than 1 pint.

The R. lung was congested and oedematous. At the root of the lung, just behind Bronchus a vertical wound of the lung was discovered, measuring about 4 inches in length and $\frac{1}{4}$ inch in breadth.

The L. lung was much congested and very oedematous a small triangular wound was found at the surface of lower lobe, laterally.

The Bronchi contained small clots, and thin m. m. was stained in both lungs.

Heart normal semi-contracted R auricle distended with red clot.

Liver 68 oz normal

Kidneys 8 oz: L - 6 oz: R. at the surface of L. kidney a few cysts were observed; the kidney showed patches of congestion the cellular tissue surrounding the R kidney contained extravasated blood.

Thorax: Extensive ecchymoses along the ribs, under the pleura on the R. side the 4. 5. 6. 7. & 8th ribs were fractured just in front of the axillary line.

On the L. side the 2. 3. 4. 5. 6. 7 & 8th ribs. at a distance of about 3 inches from the vertebral column. the 5th rib also near axillary line.

The 6th intervertebral disc was completely lacerated, the body of the 5th vertebra somewhat shattered at its base, and slightly displaced, along with the upper part of the spine, to the right.

Dr Ewart

CASE

A Dray on which patient was riding upset throwing him violently to the ground, one of the horses which had also fallen then rolled on him. On admission Jan 18. Patient was sensible, very palled, the lips were blue, great faintness was present, and skin was very cold & clammy. There was intense pain in upper dorsal region of spine, with complete loss of power and impaired sensation, in lower limbs, abdomen and chest as high as level of 1st dorsal vertebra. There was diaphragmatic breathing - Motions were passed involuntarily and the bladder was powerless.

Patient sank and died 8 hours after admission.

William St Bennett

No.	NAME	Age	Admitted	Died	Under the care of	Examination, at what time after death
20	Thomas Hindley	50	18 th Jan 77	18 th Jan	W ^h Holmes	26 hours

MORBID APPEARANCES

5 f. 8 Body well nourished, full habit. No visible trace of external injury, with the exception of a little marbling of the skin at precordial region and some bulging of the R side of thorax posteriorly.

Hair light brown

Lungs. The R. pleura contained about 2 pints of fluid blood. The L. pleura rather more than 1 pint.

The R. lung was congested and oedematous. At the root of the lung, just behind Bronchus a vertical wound of the lung was discovered, measuring about 4 inches in length and 1/2 inch in breadth.

The L. lung was much congested and very oedematous a small triangular wound was found at the surface of lower lobe, laterally.

The Bronchi contained small clots, and thin m. m. was stained in both lungs.

Heart normal semi-contracted R auricle distended with red clot.

Liver 68 oz normal.

Kidney 8 oz L - 6 oz R. at the surface of L. kidney a few cysts were observed: the kidney showed patches of congestion the cellular tissue surrounding the R kidney contained extravasated blood.

Thorax: Extensive ecchymoses along the ribs, under the pleura on the R. side the 4.5.6.7. & 8th ribs were fractured just in front of the axillary line.

On the L. side the 2.3.4.5.6.7 & 8th ribs. at a distance of about 3 inches from the vertebral column. the 5th rib also near axillary line.

The 6th intervertebral disc was completely lacerated, the body of the 5th vertebra somewhat shattered at its base, and slightly displaced, along with the upper part of the spine, to the right.

D. Ewart

No. in Reg ^t 1877	NATURE OF DISEASE
82	Fractured ribs. Haemothorax. Luxation of spine

CASE

At Dray on which patient was riding upset throwing him violently to the ground, one of the horses which had also fallen then rolled on him. On admission Jan 18 Patient was sensible, very palled, the lips were blue, great faintness was present, and skin was very cold & clammy. There was intense pain in upper dorsal region of spine, with complete loss of power and

in lower limbs, abdomen level of 1st dorsal vertebra. ratic breathing - motions rarely and the bladder was

died 8 hours after admission.

	Pyemia	1877
11	Chas. Fenwick.	P.M.
	Originated in the abortion.	
69	Albert Hall.	P.M. 25
	Myocarditis	
160	Bridget Duff	P.M. 95
	no apparent origin	
508	Emily Bannister.	P.M.
	Abscess on leg	
To 809.	Edmund Butler	Recovery.

Thomas H. Bennett

1877 p 20

No.	NAME	Age	Admitted	Died	Under the care of	Examination, at what time after death
20	Thomas Hindley	50	18 th Jan 77	18 th Jan	W ^h Holmes	26 hours

MORBID APPEARANCES

5 L. 8 Body well nourished, full habit. No visible trace of external injury, with the exception of a little marbling of the skin at precordial region and some bulging of the R side of thorax posteriorly.

Hair light brown

Lungs. The R. pleura contained about 2 pints of fluid blood. The L. pleura rather more than 1 pint.

The R. lung was congested and oedematous. At the root of the lung, just behind Bronchus a vertical wound of the lung was discovered, measuring about 4 inches in length and $\frac{1}{2}$ inch in breadth.

The L. lung was much congested and very oedematous a small triangular wound was found at the surface of lower lobe, laterally.

The Bronchi contained small clots, and thin m. m. was stained in both lungs.

Heart normal semi contracted R auricle distended with red clot.

Liver 68 oz normal

Kidneys 8 oz - L - 6 oz - R. At the surface of L. kidney a few cysts were observed: the kidney showed patches of congestion the cellular tissue surrounding the R kidney contained extravasated blood.

Thorax: Extensive ecchymoses along the ribs, under the pleura on the R. side the 4. 5. 6. 7. & 8th ribs were fractured just in front of the axillary line.

On the L. side the 2. 3. 4. 5. 6. 7. & 8th ribs. at a distance of about 3 inches from the vertebral column. the 5th rib also near axillary line.

The 6th intervertebral disc was completely lacerated, the body of the 5th vertebra somewhat shattered at its base, and slightly displaced, along with the upper part of the spine, to the right.

See Enant.

No. in Reg ^r 1877	NATURE OF DISEASE
82	Fractured ribs. Haemothorax. Luxation of spine

CASE

At Dray on which patient was riding upset throwing him violently to the ground, one of the horses which had also fallen then rolled on him.

On admission Jan 18. Patient was sensible, very palled, the lips were blue, great faintness was present, and skin was very cold & clammy.

There was intense pain in upper dorsal region of spine, with complete loss of power and impaired respiration, in lower limbs, abdomen and chest as high as level of 1st dorsal vertebra.

etc breathing - motions

and the bladder was

- 1878.
125. Fred. Ayer. P.M. 27.
Myocarditis - abrus
150. Fred. Williams. P.M.
Disease of petrous bone
324. Anne Collier
Appendicitis from P. l. origin.
Left at her own request.
522. Richard von Coppelpalle. P.M. 114
574. Eliza Patching.
Transferred to Surgeons.

died 8 hours after admission

Janus St. Bennett