

No.	NAME	Age	Admitted	Died	Under the care of	Examination, at what time after death
380	W <sup>m</sup> Rippingle Kruer	27	Nov. 17 <sup>th</sup>	Nov 22 <sup>nd</sup>	W. Dickerson	14 Hours

MORBID APPEARANCES

**General Cond<sup>n</sup>** - Fairly nourished 5'8" : Hair dark.

**Lungs etc.** *long* Congested, especially at base behind pleura natural.

**Heart** *Very prominent* 13 oz: Left. Ventricle contracted and hypertrophied: Substance of heart paler and softer than natural. Valves healthy.

**Liver** *Fatty* 4 lbs 5 oz: pale-yellow, soft, greasy. G. Bladder partly filled.

**Spleen** *long* 12 oz: soft & congested.

**Kidneys** *long* 16 oz: Large: congested: healthy.

**Alimentary C.** *Typhoid ulceration* Full of yellow putrescent faeces. Peyers Patches in lowest portion of Ileum, especially near Ileo Caecal Valve, enlarged, prominent, soft & some slightly ulcerated on surface: congested area surrounding them. The ulceration extending into submucous coat only. The large intestine was considerably congested at upper part but there was no ulceration or enlargement of solitary glands. The caecum & sigmoid glands enlarged & softened.

**Brain.** *Fluid in Ventricles.* Much serum fluid beneath Pia Mater: Lat. Ventricle distended with clear fluid: Pineal Mass. not marked: Brain substance natural.

Brain 4.3 gm

No. in Reg <sup>r</sup> 1887	NATURE OF DISEASE
1413	Typhoid.


CASE

**History.** An imperfect history was obtainable of a gradual onset of illness about a fortnight before admission characterized by headache, loss of appetite and strength with pain in the back of the neck, and a week later with much diarrhoea. He had been delirious for two nights before admission -

**or Admission.** Nov. 17. King 23. a strongly built man with congested venous face and dull heavy look: some dyspnoea. P. 102 weak, empty, no cough. Heart first sound very weak. Abdomen tender & gurgling in iliac fossa: no tympanite: one or two typhoidal maculae. He lies on his back in a state of prostration and indifference, groaning occasionally.

**Progress.** He was treated with Quin Sulph 8i. Vin. Gly. 6i. & 6 oz of Brandby increased the day before his death to 8 oz. He did not rally in any respect: After the first night which was comparatively quiet he passed each night in violent delirium which was quieter during the day. His pulse did not grow stronger or quicker (until just before his death) varying from 92 to 112. His motions were putrescent, ~~not~~ followed, on an average two a day: passed under him - He took his

by and never vomited it: he showed no signs of pain or prostration; typhoidal maculae appeared after admission - on the 25<sup>th</sup> when his temperature was at 102° 6 a dose of Sodae salicyl ʒʒ was given, but no reduction of temperature was observed: 6 hours it was 102° 8. On the evening of 28<sup>th</sup> his breathing went to 36 also 15 150: at 10 P.M. his temp. had risen to 108° 4 and he died.



A. T. Myers.

No.	NAME	Age	Admitted	Died	Under the care of	Examination, at what time after death
380	Wm. Rippling Kruer	27	Nov. 17 <sup>th</sup>	Nov. 22 <sup>nd</sup>	D <sup>r</sup> Dickerson	14 hours

MORBID APPEARANCES

*General Cond<sup>n</sup>* Fairly nourished 5'8" : Hair dark.

*Lungs etc.* Congested, especially at base behind  
*lung<sup>s</sup>* Pleurae natural.

*Heart* 13 lbs: Left. Ventricle contracted and  
*Myocardium pale* hypertrophied:  
Substance of heart paler and softer than  
natural.  
Valves healthy.

*Liver* 4 lbs 5 oz: pale-yellow, soft, greasy.  
*Fatty* G. Bladder partly filled.

*Spleen* 12 lbs: soft & congested.  
*lung<sup>s</sup>*

*Kidneys* 16 oz: Large: congested: healthy.  
*lung<sup>s</sup>*

*Alimentary C.* Full of yellow putrescent faeces.  
*Typhoid ulceration* Peyer's Patches in lower portion of Ileum  
Especially near Ileo Coecal Valve, enlarged  
prominent, soft & some slightly ulcerated on  
Surface: congested area surrounding them.  
The ulceration extending into submucous coat  
The large intestine was considerably congested  
upper part but there was no ulceration or enlargement  
of Solitary glands.  
Mesenteric glands enlarged & softened.

*Brain.* Much serum fluid beneath Pia Mater: Lat. &  
*fluid in ventricles* distributed with clear fluid: Pons & Med. not marked  
Brain substance natural.

Rey 4: Z...

No. in Reg <sup>t</sup> 188/	NATURE OF DISEASE
1413	Typhoid.

CASE

*History.* An imperfect history was obtainable of a gradual onset of illness about  
a fortnight before admission characterized by headache, loss of appetite and  
strength with pain in the back of the neck, and a week later with much  
diarrhoea. He had been delirious for two nights before admission -  
*or Admission.* Nov. 17. King 23. a strongly built man with congested venous face and dull heavy  
look: some dyspnoea - P. 102 weak, empty, no cough. Heart first sound very  
weak. *Admes* tenderness & gurgling in A. iliac fossa: no tympanite: one or two  
typhoidal maculae. He lies on his back in a state of prostration and  
indifference, groaning occasionally.

*Prognosis.* He was treated with Quin Sulph 80 ii. Vin<sup>g</sup>. q. s. and a 6 oz of Brandy increased  
the day before his death to 8 oz. He did not rally in any respect: After the first  
night which was comparatively quiet he passed each night in violent delirium,  
which was quieter during the day. His pulse did not grow stronger or quicker  
(until just before his death) varying from 92 to 112 - His nostrils were putrescent,  
light coloured, on an average two a day: passed under him - He took his  
food readily and never vomited it: he showed no signs of pain or prostration;  
a few more typhoidal maculae appeared after admission - on the 25<sup>th</sup>  
of 21<sup>st</sup> when his temperature was at 102° 6 a dose of Sodae salicif ʒss was  
given him, but no reduction of temperature was observed: 6 hours  
afterwards it was 102° 8. On the evening of 28<sup>th</sup> his breathing rose to 36  
his pulse to 150: at 10 P.M. his temp. had risen to 108° 4 and  
at 11 P.M. he died.

A. T. Myers.