

The Ovary continued
lines in diameter. This has been completely closed by
a thick layer uniting the portions of gut in which
was found to the parietes just above the pubes.
true adjoining this perforation were several points of
in the submucous and subserous cellular tissues
part of the abdominal cavity there was a quantity
fluid mixed with thin fecal matter and shreds

No 3044.

No 43. admitted 14th December 1843 under Dr. Seymour
and died on the 28th
appearances 18 hours after death. The body is exceedingly
and the countenance exhibits the cadaveric hue and
features which persons labouring under Malignant
frequently have. There is some oedema of the left lower
On opening the Thorax the left pleural cavity was
completely distended with a dark straw coloured serum
with lymph and the membrane exhibiting no traces
inflammation. The lung was compressed against the spine
into a mass resembling spleen in texture it
no air and when placed in water sunk to the bottom
lung was healthy but congested especially at the
The heart was smaller than natural but healthy
The abdominal cavity contained a quantity
fluid the lining membrane being thickened
The intestines generally were glued to one another
neighbouring organs by lymph and recently organized

Mary Joyce continued
cellular adhesions. The omentum had quite a
structure and was with difficulty recognized
firm puckered map lying underneath the liver
to it and the neighbouring organs. when cut
exhibited a gristly texture composed of firm fibres
interlacing in every direction from the interstices
1843. No in Register 2465.

No 267. Henry France At 30 admitted Oct. 12 one for
Died Dec 29 under the care of Dr. Haime
P.M. 13 hours after death. Dropsy diseased kidneys water
Body well developed and muscular. general change
Anasarca over the whole body. Countenance exte
lined with a quantity of froth around the
Mouths.

Head. all the parts within the cranium contained
more serum than natural. and the substance
of the brain itself was remarkably wet.
The cerebellum soft but without any
apparent change of structure.
Thorax. Some fluid was contained in both pleural
cavities but principally upon the left side.
The lungs were gorged with serum and
blood posteriorly and very soft in texture
The heart presented a white patch over
the right ventricle. The left ventricle consid-
erably hypertrophied unattended by any other
disease of the organ.
abdomen. The kidneys were rather above their

John Oliver continued

lines in diameter. This had been completely closed by
a thick layer uniting the portions of gut in which
was found to the parietes just above the pubes.
time adjoining this perforation were several points of
in the submucous and subserous cellular tissues
part of the abdominal cavity there was a quantity
fluid.

At 45: ad
and dia
appearan
and
features
frequ
On op
pletel
with ly
inflam
used in
no air
lung, w
The Sec
herts
ured f
The
eighbo

Many Jeyes continued
cellular adhesions. The omentum had quite a
structure and was with difficulty recognized,
firm puckered map lying underneath the liver
to it and the neighbouring organs. when cut
exhibited a gristly texture composed of firm fibres
interlacing in every direction from the intestine
which a thick white fluid might be squeezed
was healthy. The stomach contracted to one fourth
size was adherent to the neighbouring organs.
was dwindled and of firmer texture than water.
Presentery had undergone a similar change
presenting a map of schirrhous substance extending
underneath the spleen across the spine to the right.
The bladder the rectum and uterus were
in a map of ^{schirrhous} substance into which the cells
had become converted.